

Mars and frame-dragging: study for a dedicated mission

Lorenzo Iorio
INFN-Sezione di Pisa
Viale Unità di Italia 68, 70125
Bari (BA), Italy
tel. 0039 328 6128815
e-mail: lorenzo.iorio@libero.it

Abstract

In this paper we preliminarily explore the possibility of designing a dedicated satellite-based mission to measure the general relativistic gravitomagnetic Lense-Thirring effect in the gravitational field of Mars. The focus is on the systematic error induced by the multipolar expansion of the areopotential and on possible strategies to reduce it. It turns out that the major sources of bias are the Mars'equatorial radius R and the even zonal harmonics $J_\ell, \ell = 2, 4, 6, \dots$ of the areopotential. An optimal solution, in principle, consists of using two probes at high-altitudes ($a \approx 9500 - 9600$ km) and different inclinations (one probe should fly in a nearly polar orbit), and suitably combining their nodes in order to entirely cancel out the bias due to δR . The remaining uncanceled mismodelled terms due to $\delta J_\ell, \ell = 2, 4, 6, \dots$ would induce a bias $\lesssim 1\%$, according to the present-day MGS95J gravity model, over a wide range of admissible values of the inclinations. The Lense-Thirring out-of-plane shifts of the two probes would amount to about 10 cm yr^{-1} .

Keywords: Experimental tests of gravitational theories; Lunar, planetary, and deep-space probes; Mars; Gravitational fields

PACS: 04.80.Cc; 95.55.Pe; 96.30.Gc; 96.12.Fe

1 Introduction

Recent claims concerning a detection of the general relativistic gravitomagnetic Lense-Thirring effect [1] in the gravitational field of Mars with the Mars Global Surveyor probe [2, 3, 4, 5] raised interest concerning such a new Solar System scenario for testing Einsteinian gravity. In view of what happened to

the Gravity Probe B mission [6, 7], which might finally reach an accuracy¹ far from the originally expected 1% in measuring the Schiff precessions [8] of the spins of four gyroscopes carried onboard, and of the lingering controversy about the realistic accuracy reached with the terrestrial LAGEOS-LAGEOS II test of the Lense-Thirring effect [9] (and reachable with the recently approved LAGEOS-like LARES satellite, to be launched at the end of² 2009 by the Italian Space Agency [10]) we feel that attempts to scour new and unexplored routes may be of some value. Indeed, as many laboratories and methods as possible should be used to extensively test a fundamental interaction like gravity and its predictions, especially when their tests are so few and their outcomes uncertain.

In this paper we wish to fix a first stick by studying in some details the possibility of using a dedicated mission to Mars with one or more probes to measure at a reasonable level of accuracy (a few percent) the elusive Lense-Thirring effect on the longitude of the ascending node Ω of the spacecraft's orbital plane. We will concentrate here on one of the major source of systematic error, i.e. the multipolar expansion of the martian gravitational potential which induces on the node a huge noise with the same temporal signature of the relativistic signal of interest (a linear rate) in order to see what are the critical issues in view of the present-day knowledge of the Martian space environment. We will not discuss here the perturbations of non-gravitational origin which depend on the shape, the instrumentation and the orbital maneuvers of such probes. They would be strongly related to possible other tasks, more consistent with planetology, which could be fruitfully assigned to such a mission in order to enhance the possibility that it may become something more than a mere, although-hopefully-interesting, speculation; in this respect the medium-long term ambitious programs of NASA to Mars may turn out to be useful also for the purpose discussed here.

¹See on the WEB <http://einstein.stanford.edu/>

²See on the WEB http://www.esa.int/esapub/bulletin/bulletin135/bul135f_bianchi.pdf

2 The use of one nearly polar spacecraft

The Lense-Thirring effect consists of a small secular precession of the node³ Ω of the orbit of a satellite moving around a central slowly rotating mass

$$\dot{\Omega}_{\text{LT}} = \frac{2GS}{c^2 a^3 (1 - e^2)^{3/2}}, \quad (1)$$

where G is the Newtonian constant of gravitation, S is the spin angular momentum of the central body, a and e are the semimajor axis and the eccentricity, respectively, of the satellite orbit. According to the latest determinations of the global properties of Mars [11], $S = (1.92 \pm 0.01) \times 10^{32} \text{ kg m}^2 \text{ s}^{-1}$ for the red planet.

The oblateness of the central body, of mass M and equatorial radius R , makes the node to secularly precess as well according to [12]

$$\dot{\Omega}^{(\text{obl})} = \sum_{\ell=2} \dot{\Omega}_{\ell} J_{\ell}, \quad (2)$$

where the coefficients $\dot{\Omega}_{\ell}$, explicitly computed in, e.g., [13] up to degree $\ell = 20$, depend on the orbital semimajor axis, the eccentricity and the inclination i to the planet's equator; for example, for $\ell = 2$ we have

$$\dot{\Omega}_{2,2} = -\frac{3}{2} n \left(\frac{R}{a} \right)^2 \frac{\cos i}{(1 - e^2)^2}, \quad (3)$$

where $n = \sqrt{GM/a^3}$ is the probe's Keplerian mean motion. Although the multipolar expansion of the classical part of the gravitational potential is usually included in the dynamical force models used to process probes' data, such a modelling is always imperfect; thus, an aliasing trend due to the mismodelling in the even zonals J_{ℓ} and the other parameters like R and GM entering the classical node precessions may affect the recovery of the relativistic precession of interest.

2.1 The even zonal harmonics

Let us, now, focus on the systematic error $\delta\mu$ induced by the uncertainty in the even zonals δJ_{ℓ} by assuming for them the calibrated covariance sigmas of the latest Mars gravity model MGS95J [11]. We will evaluate it as

$$\delta\mu_{J_{\ell}} \leq |\dot{\Omega}_{\ell}| \delta J_{\ell}, \quad \ell = 2, 4, \dots \quad (4)$$

³Also the pericentre ω of a satellite secularly precesses under the action of the gravitomagnetic force, but we will not consider here such an effect.




Figure 1: Systematic percent errors $\delta\mu_{J_\ell}$ per degree ℓ due to the mismodelling in the even zonal harmonics δJ_ℓ , $\ell = 2, 4, 6, \dots$ for $a = 3796$ km, $e = 0.01$, $89 \text{ deg} \leq i \leq 91 \text{ deg}$ according to the calibrated covariance sigmas of the MGS95J global gravity solution up to $\ell = 20$. The Lense-Thirring effect amounts to 34 mas yr^{-1} corresponding to a shift in the out-of-plane direction of 0.62 m yr^{-1} .

Because of the presence of $\cos i$ in all the coefficients $\dot{\Omega}_\ell$ of the oblateness-induced node precessions, we will concentrate on nearly polar ($i \approx 90 \text{ deg}$) orbital configurations to minimize such a corrupting effect.

It turns out that altitudes of a few hundreds of km typical of the majority of the currently ongoing martian missions are definitely not suited for our scope, as shown by Figure 1. Note that inserting a probe into an areocentric orbit is not an easy task, so that we decided to allow for a departure of up to 1 deg from the ideal polar orbital configuration to account for unavoidable orbital injection errors. Also typical mission requirements pull the inclination some degrees apart from 90 deg: for Mars Global Surveyor (MGS) $i = 92.86 \text{ deg}$. The systematic bias per degree increases for higher degrees $\ell = 2, 4, 6, \dots$ and it turns out that only a very narrow range for i ,




Figure 2: Systematic percent errors $\delta\mu_{J_\ell}$ per degree ℓ due to the mismodelling in the even zonal harmonics δJ_ℓ , $\ell = 2, 4, 6 \dots$ for $a = 9500$ km, $e = 0.01$, $89 \text{ deg} \leq i \leq 91 \text{ deg}$ according to the calibrated covariance sigmas of the MGS95J gravity model up to $\ell = 20$. The Lense-Thirring effect amounts to 2 mas yr^{-1} corresponding to a shift in the out-of-plane direction of 0.10 m yr^{-1} .

i.e. $\Delta i \approx 10^{-3} \text{ deg}$, unlikely to obtain, might push the systematic errors below the 1% level.

By keeping a near polar geometry, much larger values of the semimajor axis, comparable with that of Phobos ($a = 9380 \text{ km}$), one of the two natural satellites of Mars, yield reasonable results, as shown by Figure 2. Indeed, now the noise decreases with high degree terms and the most serious contribution is due to the first even zonal $\ell = 2$ yielding a bias up to about 11% for $\Delta i = 1 \text{ deg}$; J_4 affects the the Lense-Thirring precessions at most at 3% level for $\Delta i = 1 \text{ deg}$. Note also that, now, a much larger departure from 90 deg ($\Delta i \approx 0.1 \text{ deg}$) would allow a further reduction of the noise ($\lesssim 1\%$). It must be noted that for such a high-altitude orbital configuration the gravitomagnetic shift in the out-of-plane direction would amount to 0.10 m yr^{-1} ;

it is certainly a small figure, but, perhaps, not too small if one considers that the average shift in the out-of-plane direction of the much lower Mars Global Surveyor is 1.6 m after about 5 yr. It poses undoubtedly challenging requirements in terms of sensitivity and overall orbit determination accuracy, but it seems not unrealistic to hope that future improvements may allow to detect such a small displacement.

Finally, let us remark that, in principle, also the temporal variations of J_2 should be accounted for; however, since at present no secular trends have been detected, such changes, mainly seasonal, annual and semi-annual [11] would not seriously impact our measurement.

2.2 The impact of R and GM

Another martian parameter which must be taken into account is the equatorial radius R along with its uncertainty. Since

$$\dot{\Omega}_{,\ell} \propto R^\ell, \quad (5)$$

the systematic error per degree induced by δR can be written as

$$\delta\mu_{J_\ell} \leq \ell \left(\frac{\delta R}{R} \right) \left| \dot{\Omega}_{,\ell} J_\ell \right|, \quad \ell = 2, 4, 6... \quad (6)$$

If we assume conservatively for it $\delta R = 6080$ m, i.e. the difference between the reference value of MGS95J [11] and the one in [14], the result is depicted in Figure 3 for $a = 9500$ km. If, instead, one takes $\delta R = 0.04$ km the errors per degree are as in Figure 4 for $a = 9500$ km. As can be noted, R is the most serious source of systematic error, even for high altitudes. Moreover, it must also be considered that it would be likely unrealistic to expect improvements in the determination of R by future areocentric missions able to push δR below terrestrial values, i.e. $1 - 0.1$ m. Thus, R will likely remain an insurmountable obstacle if only one probe is to be used.

Concerning GM , it turns out that the impact of its mismodelling is of no concern being $\lesssim 1\%$.

2.3 Summary

In summary, in this Section we investigated a nearly polar, high-altitude Phobos-like orbit and noted that the Lense-Thirring effect amounts to a shift of 0.10 m yr^{-1} in the out-of-plane direction. Concerning the systematic errors, the most crucial source of aliasing is the radius R ($\delta\mu \approx 10000\%$ for $\delta R = 0.04$ km) and, to a lesser extent, the first even zonal harmonic




Figure 3: Systematic percent errors $\delta\mu_{J_\ell}$ per degree ℓ due to the uncertainty in Mars' radius R , assumed to be $\delta R = 6080$ m, for $a = 9500$ km, $e = 0.01$, $89 \text{ deg} \leq i \leq 91 \text{ deg}$. The errors for $\ell = 4, 6$ are as large as 6000% and 500%, respectively.




Figure 4: Systematic percent errors $\delta\mu_{J_\ell}$ per degree ℓ due to the uncertainty in Mars' radius R , assumed to be $\delta R = 0.04 \text{ km}$, for $a = 9500 \text{ km}$, $e = 0.01$, $89 \text{ deg} \leq i \leq 91 \text{ deg}$. The errors for $\ell = 4, 6$ are as large as 40% and 5%, respectively.

J_2 of the areopotential ($\delta\mu \leq 11\%$). To reduce them the probe should be inserted into an orbit with an inclination close to 90 deg within 10^{-3} deg or less. In conjunction with such a very tight constraint, it must also be hoped that future missions to Mars will improve our knowledge of the fundamental parameters of the red planet to a sufficient extent to allow for larger departures from the ideal polar geometry: after all, the gravity solution MGS95J represents an improvement of one order of magnitude with respect to the previous MGS75D model [15]. However, this may be valid for the spherical harmonic coefficients of the areopotential, not for the radius of Mars for which an uncertainty of the order of the meter would not be enough. We must conclude that the use of only one probe for measuring its node Lense-Thirring precession is unfeasible.

3 Including Phobos and Deimos

A possible solution to reduce the systematic errors would be to consider a linear combination of the nodes of the proposed probe and of the two natural satellites of Mars, i.e. Phobos ($a = 9380$ km, $i = 1.075$ deg, $e = 0.0151$) and Deimos ($a = 23460$ km, $i = 1.793$ deg, $e = 0.0002$), suitably designed to cancel out the impact of just δJ_2 and $\delta R/R$, according to an approach put forth for the first time in the context of the terrestrial LAGEOS-LAGEOS II test [16]. A possible combination is

$$\delta\dot{\Omega}^{\text{Deimos}} + c_1\delta\dot{\Omega}^{\text{Probe}} + c_2\delta\dot{\Omega}^{\text{Phobos}}, \quad (7)$$

with $\delta\dot{\Omega}$ denoting some sort of Observed-minus-Calculated ($O - C$) quantity accounting for any unmodelled/mismodelled features of motion and extracted from the data processing with full dynamical force models. By purposely leaving the gravitomagnetic force unmodelled, it can be written down as

$$\delta\dot{\Omega} = \mu X_{\text{LT}} + \dot{\Omega}_{.2}\delta J_2 + \dot{\Omega}_{.R} \left(\frac{\delta R}{R} \right) + \Delta : \quad (8)$$

here $\mu = 1$ in general relativity, the coefficient $\dot{\Omega}_{.R}$ is defined as

$$\dot{\Omega}_{.R} = R \frac{\partial \dot{\Omega}^{(\text{obl})}}{\partial R} = \sum_{\ell=2} \ell \dot{\Omega}_{.\ell} J_{\ell}, \quad (9)$$

and Δ represents all other remaining mismodelled/unmodelled effects acting on the node (e.g. $\delta J_4, \delta J_6, \dots; \delta GM$). By solving for μ one gets

$$\begin{cases} c_1 = \frac{\dot{\Omega}_{2.2}^{\text{Phobos}} \dot{\Omega}_R^{\text{Deimos}} - \dot{\Omega}_{2.2}^{\text{Deimos}} \dot{\Omega}_R^{\text{Phobos}}}{\dot{\Omega}_{2.2}^{\text{Probe}} \dot{\Omega}_R^{\text{Phobos}} - \dot{\Omega}_{2.2}^{\text{Phobos}} \dot{\Omega}_R^{\text{Probe}}}, \\ c_2 = \frac{\dot{\Omega}_{2.2}^{\text{Deimos}} \dot{\Omega}_R^{\text{Probe}} - \dot{\Omega}_{2.2}^{\text{Probe}} \dot{\Omega}_R^{\text{Deimos}}}{\dot{\Omega}_{2.2}^{\text{Probe}} \dot{\Omega}_R^{\text{Phobos}} - \dot{\Omega}_{2.2}^{\text{Phobos}} \dot{\Omega}_R^{\text{Probe}}}. \end{cases} \quad (10)$$

Note that c_1 , contrary to c_2 , is not defined for $i = 90$ deg because $\dot{\Omega}_\ell^{\text{Probe}}$, which are proportional to $\cos i$, vanish for all ℓ at $i = 90$ deg. The result, in principle, would be satisfactory. Indeed, now, the systematic bias of $J_4, J_6, J_8 \dots$ on the combination of eq. (7) turns out to be $\delta\mu \lesssim 6\%$ in $89 \text{ deg} \leq i \leq 89.9 \text{ deg}$; the major contribution is due to J_4 . It is an acceptable result, especially, in view of the fact that the admissible range for the inclination amounts, now, to about 1 deg and that eq. (7) is, by construction, immune to the uncertainty in the martian radius; indeed, although it is likely that many physical properties and parameters like, e.g., the even zonals, of the red planets will be determined with increasing accuracy by the many ongoing and planned missions, it is unlikely that the radius will be known to a sufficient accuracy to change at an acceptable level the bias induced by it (see Figure 3 and Figure 4).

The combination of eq. (7) is also able to remove the large part of the effect of the mismodelling in GM

$$\delta\mu_{GM} \leq \frac{1}{2} \left(\frac{\delta GM}{GM} \right) \sum_{\ell=2}^{20} \left| \dot{\Omega}_\ell J_\ell \right|. \quad (11)$$

Indeed, the main contribution to eq. (11) is due to J_2 , which is canceled out by eq. (7) with the coefficients of eq. (10). The other terms in eq. (11), not canceled by the combination of eq. (7), yield negligible errors, well below 1%.

However, it must be stressed that the node of Phobos undergoes secular precessions due to other perturbations of gravitational origin [17] (non-sphericity of Phobos itself, Mars tidal bulge and nutation) which have not been considered here and that would affect the combination of eq. (7). It turns out that the most relevant one is that due to the spherical harmonic coefficients c_{20} and c_{22} of Phobos itself [18] whose induced secular precession amounts nominally to 200 km over 3 yr [17]. The nutation perturbation nominally amounts to 0.3 km over 3 yr [17]. The tidally induced precession, parameterized in terms of the martian Love number k_2 would amount nominally 0.06 km after 3 yr [17], but being $k_2 = 0.152 \pm 0.009$ [11], its modelling

would left a $\approx 1 \text{ m yr}^{-1}$ mismodelled trend. The Phobos gravitomagnetic shift is 0.1 m yr^{-1} .

In terms of the detectability of the Lense-Thirring signal with the combination of eq. (7), undoubtedly, a major drawback of the strategy of including Phobos and Deimos is the poor accuracy with which their orbits can be reconstructed with respect to their Lense-Thirring signal. Indeed, while their gravitomagnetic shifts are of the order of $1 - 10 \text{ cm yr}^{-1}$, the latest NASA martian ephemerides⁴ for them yield an accuracy of $1 - 10 \text{ km}$ in the radial, transverse and out-of-plane orbital components [19]. Thus, according to the present-day level of accuracy including the natural satellites of Mars in the combination of eq. (7) would introduce a noise which would overwhelm the relativistic trend of interest.

4 Two dedicated probes

In principle, the linear combination approach could be followed by using the nodes of other two spacecrafts⁵, although sending to Mars three new probes would increase the costs and the difficulties of such a demanding mission.

4.1 The supplementary orbit configuration

It is interesting to consider a scenario involving only two probes, named P1 and P2, in a nearly polar counter-orbiting configuration, proposed for the first time by Van Patten and Everitt [20] in the framework of the attempts to design a suitable terrestrial mission, and later generalized by Ciufolini [21] for other inclinations. By assuming for, say, P1 $a_1 = 9500 \text{ km}$, $i_1 = 89 \text{ deg}$, $e_1 = 0.01$, it turns out that the sum of their nodes would be an observable relatively insensitive to departures from the ideal configuration for P2, i.e. $a_2 = 9500 \text{ km}$, $i_2 = 91 \text{ deg}$, $e_2 = 0.01$, at least in regard to the mismodelling in the even zonals. Indeed, it turns out that for $a_2 = 9600 \text{ km}$, $90.05 \text{ deg} \leq i_2 \leq 92 \text{ deg}$, $e_2 = 0.03$ the bias due to J_2 is up to 7%, while the other even zonals would have an impact of the order of 1% or less. Unfortunately, the uncertainty in the martian radius makes such constraints much more stringent. Indeed, $\delta\mu_R \approx 10^5\%$ for $\delta R = 6080 \text{ m}$ and $\delta\mu_R \approx 4000\%$ for $\delta R = 0.04 \text{ km}$. Also in this case, the improvements in determining R that could realistically be expected from future missions to the red planets may be not sufficient.

⁴See on the WEB http://ssd.jpl.nasa.gov/?sat_ephem.

⁵For a polar geometry the Lense-Thirring precession of the pericentre ω of a spacecraft vanishes being proportional to $\cos i$.

4.2 The linear combination approach

An approach that could be fruitfully followed within the framework of the linear combination strategy with two probes P1 and P2, especially in view of likely future improvements in the even zonal harmonics of the areopotential, consists in designing a combination which, by construction, entirely cancels out the bias due to the uncertainty in R

$$\delta\mu_R \leq \left| \dot{\Omega}_{\cdot R}^{\text{P2}} + k_1 \dot{\Omega}_{\cdot R}^{\text{P1}} \right| \left(\frac{\delta R}{R} \right), \quad (12)$$

being, instead, affected by $\delta J_2, \delta J_4, \delta J_6$. It is analogous to the three-nodes combination of eq. (7), with

$$k_1 = -\frac{\dot{\Omega}_{\cdot R}^{\text{P2}}}{\dot{\Omega}_{\cdot R}^{\text{P1}}}. \quad (13)$$

Indeed, as can be noted in Figure 5, $\delta\mu_{J_\ell}$ would be very good ($\lesssim 1\%$), according to the present-day MGS95J model, provided that inclinations $i \ll 90$ deg are adopted for the second probe. It turns out that the most relevant even zonals would be J_4 and J_6 with a bias of the order of $\approx 1 - 0.5\%$. With eq. (13) an inclination range as large as ten degrees would be admissible: it is really a great advantage with respect to the other cases examined so far.

5 Conclusions

In this paper in examining the possibility of designing a dedicated mission to measure the Lense-Thirring effect in the gravitational field of Mars by analyzing the secular precessions of the node of one or more spacecrafts we focussed on the main systematic errors of gravitational origin and strategies to reduce them.

The main source of systematic bias is the mismodelling in some of the parameters entering the multipolar expansion of the Newtonian part of the areopotential, especially the Mars' equatorial radius R and, to a lesser but non-negligible extent, the even zonal harmonics $J_\ell, \ell = 2, 4, 6, \dots$. Since the resulting aliasing node precessions are proportional to $R^\ell a^{-(3/2+\ell)} \cos i$, a high-altitude ($a \approx 9500$ km), polar ($i = 90$ deg) probe would be, in principle, an optimal solution, but the consideration of the unavoidable orbit injection errors showed that such an option is unfeasible because it would require too stringent constraints on the departures from the ideal case $i = 90$ deg. It could be possible, in principle, to suitably combine the nodes of such a probe




Figure 5: Systematic errors $\delta\mu_{J_\ell}$ per degree due to the mismodelling $\delta J_2, \delta J_4, \delta J_6, \dots$ (MGS95J model) in the uncanceled even zonal harmonics in the combination of the nodes of probe P1 ($a_1 = 9500 \text{ km}$, $i_1 = 89 \text{ deg}$, $e_1 = 0.01$) and probe P2 ($a_2 = 9600 \text{ km}$, $60 \text{ deg} \leq i_2 \leq 89 \text{ deg}$, $e_2 = 0.03$) with eq. (13) which cancels out entirely the bias due to δR .

with those of the natural satellites of Mars, Phobos and Deimos. However, the Lense-Thirring shifts of such bodies ($1 - 10 \text{ cm yr}^{-1}$) are orders of magnitude too small with respect to the present accuracy in reconstructing their orbits ($1 - 10 \text{ km}$).

A viable option consists in using two probes P1 and P2 at high-altitudes ($a_1 = 9500 \text{ km}$, $a_2 = 9600 \text{ km}$) and different inclinations ($i_1 \approx 90 \text{ deg}$, $60 \text{ deg} \leq i_2 \leq 89 \text{ deg}$), and combining their nodes so as to entirely cancel out the bias due to R : the resulting bias due to $\delta J_2, \delta J_4, \delta J_6 \dots$ would be $\lesssim 1\%$, according to the present-day MGS95J gravity model, over a range of values for the inclination as large as ten degrees.

A major challenge would certainly be reaching a satisfactory accuracy in reconstructing the orbits of such probes whose Lense-Thirring out-of-plane shifts would amount to about 10 cm yr^{-1} ; for example, the spacecraft should remain operative for many years, without any failure in the communication with the Earth, and it is likely that one or more landers would be required as well.

References

- [1] Lense, J., Thirring, H.: Über den Einfluss der Eigenrotation der Zentralkörper auf die Bewegung der Planeten und Monde nach der Einsteinschen Gravitationstheorie. *Phys. Z.* 19, 156-163 (1918)
- [2] Iorio, L.: COMMENTS, REPLIES AND NOTES: A note on the evidence of the gravitomagnetic field of Mars. *Class. Quantum Grav.* 23, 5451-5454 (2006)
- [3] Iorio, L.: Is It Possible to Measure the Lense-Thirring Effect in the Gravitational Fields of the Sun and of Mars? In: Iorio, L. (ed.) *The Measurement of Gravitomagnetism: a Challenging Enterprise*, pp. 177-187. NOVA, Hauppauge (NY) (2007)
- [4] Krogh, K.: COMMENTS, REPLIES AND NOTES: Comment on 'Evidence of the gravitomagnetic field of Mars. *Class. Quantum Grav.* 24, 5709-5715 (2007)
- [5] Iorio, L.: Reply to "Comment on 'Evidence of the gravitomagnetic field of Mars'", by Kris Krogh. *J. Gravit. Phys.*, at press, arXiv:gr-qc/0701146v4 (2008)
- [6] Everitt, C.W.F.: The Gyroscope Experiment I. General Description and Analysis of Gyroscope Performance. In: Bertotti, B. (ed.) *Proc.*

- Int. School Phys. “Enrico Fermi” Course LVI, pp. 331-360. New Academic Press, New York (1974)
- [7] Everitt, C.W.F., et al.: Gravity Probe B: Countdown to Launch. In: Lämmerzahl, C., Everitt, C.W.F., Hehl, F.W. (eds.) Gyros, Clocks, Interferometers...: Testing Relativistic Gravity in Space, pp. 52-82. Springer, Berlin (2001)
 - [8] Schiff, L.: Possible New Experimental Test of General Relativity Theory. Phys. Rev. Lett. 4, 215-217 (1960)
 - [9] Iorio, L.: An assessment of the measurement of the Lense Thirring effect in the Earth gravity field, in reply to: On the measurement of the Lense Thirring effect using the nodes of the LAGEOS satellites, in reply to On the reliability of the so far performed tests for measuring the Lense Thirring effect with the LAGEOS satellites by L. Iorio, by I. Ciufolini and E. Pavlis. Planet. Space Sci. 55, 503-511 (2007)
 - [10] Iorio, L.: Advances in the measurement of the Lense-Thirring effect with Satellite Laser Ranging in the gravitational field of the Earth. In: Koslovskiy, V.V. (ed.) Progress in Laser and Electro-Optics Research. NOVA, Hauppauge (NY). arXiv:0808.0658v1 [gr-qc]. (2008)
 - [11] Konolpiv, A.S., et al.: A global solution for the Mars static and seasonal gravity, Mars orientation, Phobos and Deimos masses, and Mars ephemeris. Icarus 182, 23-50 (2006)
 - [12] Kaula, W.M.: Theory of Satellite Geodesy. Blaisdell, Waltham (1966)
 - [13] Iorio, L.: Appendix B The Classical Orbital Precessions. In: Iorio, L. (ed.) The Measurement of Gravitomagnetism: a Challenging Enterprise, pp. 239-245. NOVA, Hauppauge (NY) (2007)
 - [14] Yoder, C.F.: Astrometric and Geodetic Properties of Earth and the Solar System. In: Ahrens, T.J. (ed.) Global earth physics a handbook of physical constants, Table 6. AGU reference shelf Series, vol. no. 1 (1995)
 - [15] Yuan, D.N., et al.: Gravity field of Mars: A 75th Degree and Order Model. J. Geophys. Res. 106 23377-23402 (2001)

- [16] Ciufolini, I.: On a new method to measure the gravitomagnetic field using two orbiting satellites. *N. Cim. A* 109, 1709-1720 (1996)
- [17] Lainey, V., Dehant, V., Pätzold, M.: First numerical ephemerides of the Martian moons. *Astron. Astrophys.* 465, 1075-1084 (2007)
- [18] Borderies, N., Yoder, C.F.: Phobos' gravity field and its influence on its orbit and physical librations. *Astron. Astrophys.* 233, 235-251 (1990)
- [19] Jacobson, R.A., Rush, B.: Ephemerides of the Martian Satellites - MAR063, JPL IOM 343R-06-004 (2006)
- [20] Van Patten, R.A., Everitt, C.W.F.: Possible Experiment with Two Counter-Orbiting Drag-Free Satellites to Obtain a New Test of Einstein's General Theory of Relativity and Improved Measurements in Geodesy. *Phys. Rev. Lett.* 36, 629-632 (1976)
- [21] Ciufolini, I.: Measurement of the Lense-Thirring drag on high-altitude, laser-ranged artificial satellites. *Phys. Rev. Lett.* 56, 278-281 (1986)